

Monday, March 23

This demonstration features the estimation of Poisson and logistic regression models, and the interpretation of rate and odds ratios.


Impact of Pesticides on Skylark Reproductivity

During the four summers from 1992 to 1995 researchers from the National Environmental Research Institute in the Ministry of Environment and Energy in Denmark conducted a study to examine how pesticide use impacts skylark reproduction in barley fields.¹ The study used a fractional factorial design in which each year two of four fields were sprayed with pesticides while the other two fields were not.² Which fields were sprayed was alternated so that a field was sprayed every other year. The number of fledgling skylarks produced in each field each year was recorded. The data are in the `skylark` data frame from the `trtools` package. The data are plotted below.

```
library(trtools)
library(ggplot2)
p <- ggplot(skylark, aes(x = spray, y = count)) +
  geom_point() + facet_grid(field ~ year) + theme_minimal() +
  labs(x = "Field Sprayed?", y = "Number of Skylark Fledglings")
plot(p)
```


¹Odderskær, P., Prang, A., Eknegaard, N., & Andersen, P. N. (1997). Skylark reproduction in pesticide treated fields (Comparative studies of *Alauda arvensis* breeding performance in sprayed and unsprayed barley fields). *Bekæmpelsesmiddelforskning fra Miljøstyrelsen*, 32, National Environmental Research Institute, Ministry of the Environment and Energy, Denmark: Danish Environmental Protection Agency.

²A [fractional factorial design](#) is a design in which observations are made at only a subset of the possible combinations of levels of two or more factors. Such designs are quite economical but can preclude the estimation of interactions. This does not mean that such interactions are not present, but rather that if they are they are confounded with the main effects. For this particular design it is only possible to fully estimate a model with “main effects” for each of the three factors. Ideally fractional factorial designs are used when interactions are negligible.




Here is another way to visualize the data that flips the axes and “combines” the field and year variables when specifying facets.

```
p <- ggplot(skylark, aes(x = count, y = spray)) +
  geom_point() + facet_grid(field + year ~ .) + theme_minimal() +
  labs(y = "Field Sprayed?", x = "Number of Skylark Fledglings")
plot(p)
```



We can use color for field to “condense” the plot a bit.

```
p <- ggplot(skylark, aes(x = count, y = spray, color = field)) +
  geom_point() + facet_grid(year ~ .) + theme_minimal() +
  labs(y = "Field Sprayed?", x = "Number of Skylark Fledglings", color = "Field")
plot(p)
```



The plots clearly shows the incomplete nature of the fractional factorial design. In any given year, a field either was or was not sprayed. The objective is to investigate the effect of spraying on the number of fledglings while controlling for the effects of year and field.

1. Estimate a Poisson regression model for the number of skylark fledglings as your response variable that will reproduce the following results.

```
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	3.430943	0.13262	25.86999	1.450e-147	3.16352	3.68367
sprayyes	-0.456126	0.09385	-4.86011	1.173e-06	-0.64141	-0.27324
fieldKr	0.049089	0.12672	0.38738	6.985e-01	-0.19929	0.29806
fieldKu	0.004964	0.12800	0.03879	9.691e-01	-0.24611	0.25625
fieldRd	-0.179048	0.13417	-1.33452	1.820e-01	-0.44342	0.08326
year1993	0.462623	0.13064	3.54108	3.985e-04	0.20868	0.72149
year1994	0.060018	0.14149	0.42420	6.714e-01	-0.21735	0.33816
year1995	0.327281	0.13411	2.44041	1.467e-02	0.06596	0.59240

Note that here `m` is a model object created using the `glm` function.

Solution: The results can be replicated as follows. Note that the output above indicates that only the “main effects” of spray, field, and year were specified. We can see that there are indicator variables for spray, field, and year, but no interaction terms.

```
m <- glm(count ~ spray + field + year, family = poisson, data = skylark)
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	3.430943	0.13262	25.86999	1.450e-147	3.16352	3.68367
sprayyes	-0.456126	0.09385	-4.86011	1.173e-06	-0.64141	-0.27324
fieldKr	0.049089	0.12672	0.38738	6.985e-01	-0.19929	0.29806
fieldKu	0.004964	0.12800	0.03879	9.691e-01	-0.24611	0.25625
fieldRd	-0.179048	0.13417	-1.33452	1.820e-01	-0.44342	0.08326
year1993	0.462623	0.13064	3.54108	3.985e-04	0.20868	0.72149
year1994	0.060018	0.14149	0.42420	6.714e-01	-0.21735	0.33816
year1995	0.327281	0.13411	2.44041	1.467e-02	0.06596	0.59240

There is no offset variable here. We will assume the fields were all of the same size. But if they were not and we knew the area of each field (in a variable called `area` for example) we might use that as an offset by specifying the model as follows.

```
m <- glm(count ~ offset(log(area)) + spray + field + year,
family = poisson, data = skylark)
```

Then we would be modeling the expected number of fledglings per unit area (e.g., number of fledglings

per square square meter).

2. What is the estimated rate ratio for the effect of spraying? How can this be interpreted?

Solution: We can estimate this rate ratio several ways. Note that since there is no interaction involving `spray` the field and year does not matter.

```
trtools::contrast(m, tf = exp,
  a = list(spray = "yes", field = "Ke", year = "1992"),
  b = list(spray = "no", field = "Ke", year = "1992"))

estimate lower upper
0.6337 0.5273 0.7617
```

We can interpret this estimated rate ratio as showing that the expected number of fledglings in a sprayed field is about 0.63 times that of a field that is not sprayed. We can also say that the expected number of fledglings in a sprayed field is about 37% less than that in a field that is not sprayed. We can “flip” the rate ratio as follows.

```
trtools::contrast(m, tf = exp,
  a = list(spray = "no", field = "Ke", year = "1992"),
  b = list(spray = "yes", field = "Ke", year = "1992"))

estimate lower upper
1.578 1.313 1.897
```

We can interpret this estimated rate ratio as showing that the expected number of fledglings in a field that is not sprayed is about 1.58 times that of a field that is sprayed, or that the number of fledglings in a field that is not sprayed is about 58% higher than a field that is sprayed.

To estimate the rate ratio using the `emmeans` package we need to use the `emmeans` function to produce the estimated expected counts and then the `pairs` function to produce rate ratios. Note that using `~spray|field*year` will allow us to produce a rate ratio for each combination of field and year.

```
library(emmeans)
pairs(emmeans(m, ~spray|field*year, type = "response"), infer = TRUE)
```

```
field = Ke, year = 1992:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes 1.58 0.148 Inf 1.31 1.9 1 4.860 <0.0001
```

```
field = Kr, year = 1992:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes 1.58 0.148 Inf 1.31 1.9 1 4.860 <0.0001
```

```
field = Ku, year = 1992:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes 1.58 0.148 Inf 1.31 1.9 1 4.860 <0.0001
```

```
field = Rd, year = 1992:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes 1.58 0.148 Inf 1.31 1.9 1 4.860 <0.0001
```

```
field = Ke, year = 1993:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes 1.58 0.148 Inf 1.31 1.9 1 4.860 <0.0001
```

```
field = Kr, year = 1993:
contrast ratio SE df asymp.LCL asymp.UCL null z.ratio p.value
```

```

no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Ku, year = 1993:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Rd, year = 1993:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Ke, year = 1994:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Kr, year = 1994:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Ku, year = 1994:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Rd, year = 1994:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Ke, year = 1995:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Kr, year = 1995:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Ku, year = 1995:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

field = Rd, year = 1995:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1      4.860 <0.0001

```

Confidence level used: 0.95

Intervals are back-transformed from the log scale

Tests are performed on the log scale

Note that by default this estimates the rate ratio for the expected number of fledglings in a field that is not sprayed to that of a field that is sprayed. To “flip” the rate ratio from the default include the option `reverse = TRUE` as follows.

```

pairs(emmeans(m, ~spray|field*year, type = "response"),
      infer = TRUE, reverse = TRUE)

```

```

field = Ke, year = 1992:
contrast ratio  SE  df  asymp.LCL  asymp.UCL  null  z.ratio  p.value

```

```

yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Kr, year = 1992:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ku, year = 1992:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Rd, year = 1992:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ke, year = 1993:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Kr, year = 1993:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ku, year = 1993:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Rd, year = 1993:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ke, year = 1994:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Kr, year = 1994:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ku, year = 1994:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Rd, year = 1994:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Ke, year = 1995:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

field = Kr, year = 1995:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001

```

```
field = Ku, year = 1995:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
```

```
field = Rd, year = 1995:
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
```

Confidence level used: 0.95
Intervals are back-transformed from the log scale
Tests are performed on the log scale

We can make a more compact display by using `rbind`, but there we need to include `adjust = "none"` to match these results because when you use `rbind`, `pairs` will assume you want to do family-wise error rate adjustments.

```
rbind(pairs(emmeans(m, ~spray|field*year, type = "response"),
infer = TRUE, reverse = TRUE), adjust = "none")
```

```
field year contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
Ke    1992 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Kr    1992 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ku    1992 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Rd    1992 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ke    1993 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Kr    1993 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ku    1993 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Rd    1993 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ke    1994 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Kr    1994 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ku    1994 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Rd    1994 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ke    1995 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Kr    1995 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Ku    1995 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
Rd    1995 yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
```

Confidence level used: 0.95
Intervals are back-transformed from the log scale
Tests are performed on the log scale

And, you can “average” over field and year by leaving them off. This produces the same result since the rate ratio is the same.

```
pairs(emmeans(m, ~spray, type = "response"), infer = TRUE)
```

```
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
no / yes  1.58 0.148 Inf      1.31      1.9      1  4.860 <0.0001
```

Results are averaged over the levels of: field, year
Confidence level used: 0.95
Intervals are back-transformed from the log scale
Tests are performed on the log scale

```
pairs(emmeans(m, ~spray, type = "response"), infer = TRUE, reverse = TRUE)
```

```
contrast ratio      SE df asymp.LCL asymp.UCL null z.ratio p.value
```

```
yes / no 0.634 0.0595 Inf      0.527      0.762      1 -4.860 <0.0001
```

Results are averaged over the levels of: field, year
Confidence level used: 0.95
Intervals are back-transformed from the log scale
Tests are performed on the log scale

It is important to note, however, that what you are averaging is *not* the rate ratio, but rather than log of the rate ratio, which is then being exponentiated. But that does not matter here since the rate ratio is the same for every field and year.

Finally the estimated rate ratio can also be found from the parameter estimates. This may not be possible for models with interactions, depending on the parameterization, but it does work here.

```
exp(cbind(coef(m), confint(m)))
```

```
                2.5 %  97.5 %  
(Intercept) 30.9058 23.6537 39.7921  
sprayyes      0.6337 0.5265 0.7609  
fieldKr       1.0503 0.8193 1.3472  
fieldKu       1.0050 0.7818 1.2921  
fieldRd       0.8361 0.6418 1.0868  
year1993     1.5882 1.2321 2.0575  
year1994     1.0619 0.8046 1.4024  
year1995     1.3872 1.0682 1.8083
```

The confidence interval is slightly different here. This is because `confint` uses what is called a profile likelihood confidence interval whereas `contrast` and functions in the `emmeans` package use what are called Wald confidence intervals.

3. What is the estimated expected number of fledglings for each condition?

Solution: This can be done several ways. For a given field and year, for example, we can estimate the expected count for field that are sprayed and not sprayed.

```
trtools::contrast(m, tf = exp,  
  a = list(spray = c("no", "yes"), field = "Ke", year = "1992"),  
  cnames = c("no spray", "spray"))
```

```
      estimate lower upper  
no spray  30.91 23.83 40.08  
spray     19.59 15.07 25.45
```

But by using `emmeans` we can easily get the estimated expected counts for all combinations of the three factors.

```
emmeans(m, ~spray|field*year, type = "response")
```

```
field = Ke, year = 1992:  
spray rate  SE  df asymp.LCL asymp.UCL  
no    30.9 4.10 Inf      23.8      40.1  
yes   19.6 2.62 Inf      15.1      25.4
```

```
field = Kr, year = 1992:  
spray rate  SE  df asymp.LCL asymp.UCL  
no    32.5 4.31 Inf      25.0      42.1  
yes   20.6 2.97 Inf      15.5      27.3
```

```
field = Ku, year = 1992:
```

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	31.1	4.16	Inf	23.9	40.4
yes	19.7	2.87	Inf	14.8	26.2

field = Rd, year = 1992:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	25.8	3.58	Inf	19.7	33.9
yes	16.4	2.28	Inf	12.5	21.5

field = Ke, year = 1993:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	49.1	6.11	Inf	38.5	62.6
yes	31.1	3.94	Inf	24.3	39.9

field = Kr, year = 1993:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	51.6	5.59	Inf	41.7	63.8
yes	32.7	4.04	Inf	25.6	41.6

field = Ku, year = 1993:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	49.3	5.42	Inf	39.8	61.2
yes	31.3	3.90	Inf	24.5	39.9

field = Rd, year = 1993:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	41.0	5.37	Inf	31.8	53.0
yes	26.0	3.45	Inf	20.1	33.7

field = Ke, year = 1994:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	32.8	4.28	Inf	25.4	42.4
yes	20.8	2.73	Inf	16.1	26.9

field = Kr, year = 1994:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	34.5	4.50	Inf	26.7	44.5
yes	21.8	3.11	Inf	16.5	28.9

field = Ku, year = 1994:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	33.0	4.34	Inf	25.5	42.7
yes	20.9	3.00	Inf	15.8	27.7

field = Rd, year = 1994:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	27.4	3.74	Inf	21.0	35.8
yes	17.4	2.39	Inf	13.3	22.8

field = Ke, year = 1995:

spray rate	SE	df	asyp.LCL	asyp.UCL	
no	42.9	5.49	Inf	33.4	55.1
yes	27.2	3.54	Inf	21.1	35.1

```
field = Kr, year = 1995:
  spray rate  SE  df asymp.LCL asymp.UCL
no    45.0 5.07 Inf      36.1    56.1
yes   28.5 3.63 Inf      22.2    36.6
```

```
field = Ku, year = 1995:
  spray rate  SE  df asymp.LCL asymp.UCL
no    43.1 4.91 Inf      34.5    53.9
yes   27.3 3.51 Inf      21.2    35.1
```

```
field = Rd, year = 1995:
  spray rate  SE  df asymp.LCL asymp.UCL
no    35.8 4.81 Inf      27.6    46.6
yes   22.7 3.09 Inf      17.4    29.7
```

Confidence level used: 0.95

Intervals are back-transformed from the log scale

The output will be organized a little differently if we use `~spray*field*year`.

```
emmeans(m, ~spray*field*year, type = "response")
```

```
spray field year rate  SE  df asymp.LCL asymp.UCL
no    Ke   1992 30.9 4.10 Inf      23.8    40.1
yes   Ke   1992 19.6 2.62 Inf      15.1    25.4
no    Kr   1992 32.5 4.31 Inf      25.0    42.1
yes   Kr   1992 20.6 2.97 Inf      15.5    27.3
no    Ku   1992 31.1 4.16 Inf      23.9    40.4
yes   Ku   1992 19.7 2.87 Inf      14.8    26.2
no    Rd   1992 25.8 3.58 Inf      19.7    33.9
yes   Rd   1992 16.4 2.28 Inf      12.5    21.5
no    Ke   1993 49.1 6.11 Inf      38.5    62.6
yes   Ke   1993 31.1 3.94 Inf      24.3    39.9
no    Kr   1993 51.6 5.59 Inf      41.7    63.8
yes   Kr   1993 32.7 4.04 Inf      25.6    41.6
no    Ku   1993 49.3 5.42 Inf      39.8    61.2
yes   Ku   1993 31.3 3.90 Inf      24.5    39.9
no    Rd   1993 41.0 5.37 Inf      31.8    53.0
yes   Rd   1993 26.0 3.45 Inf      20.1    33.7
no    Ke   1994 32.8 4.28 Inf      25.4    42.4
yes   Ke   1994 20.8 2.73 Inf      16.1    26.9
no    Kr   1994 34.5 4.50 Inf      26.7    44.5
yes   Kr   1994 21.8 3.11 Inf      16.5    28.9
no    Ku   1994 33.0 4.34 Inf      25.5    42.7
yes   Ku   1994 20.9 3.00 Inf      15.8    27.7
no    Rd   1994 27.4 3.74 Inf      21.0    35.8
yes   Rd   1994 17.4 2.39 Inf      13.3    22.8
no    Ke   1995 42.9 5.49 Inf      33.4    55.1
yes   Ke   1995 27.2 3.54 Inf      21.1    35.1
no    Kr   1995 45.0 5.07 Inf      36.1    56.1
yes   Kr   1995 28.5 3.63 Inf      22.2    36.6
no    Ku   1995 43.1 4.91 Inf      34.5    53.9
yes   Ku   1995 27.3 3.51 Inf      21.2    35.1
no    Rd   1995 35.8 4.81 Inf      27.6    46.6
yes   Rd   1995 22.7 3.09 Inf      17.4    29.7
```

Confidence level used: 0.95

Intervals are back-transformed from the log scale


Note that the arguments `tf = exp` and `type = "response"` are necessary when using `contrast` and `emmeans`, respectively, so that that we are estimating the expected response rather than the log of the expected response. Another approach is to use the `glmint` function from the `trtools` package.

```
d <- expand.grid(spray = c("yes", "no"), field = c("Ke", "Kr", "Ku", "Rd"),
  year = c("1992", "1993", "1994", "1995"))
cbind(d, trtools::glmint(m, newdata = d))
```

	spray	field	year	fit	low	upp
1	yes	Ke	1992	19.59	15.07	25.45
2	no	Ke	1992	30.91	23.83	40.08
3	yes	Kr	1992	20.57	15.50	27.31
4	no	Kr	1992	32.46	25.02	42.11
5	yes	Ku	1992	19.68	14.80	26.18
6	no	Ku	1992	31.06	23.88	40.39
7	yes	Rd	1992	16.38	12.46	21.52
8	no	Rd	1992	25.84	19.69	33.90
9	yes	Ke	1993	31.11	24.27	39.87
10	no	Ke	1993	49.09	38.46	62.65
11	yes	Kr	1993	32.67	25.64	41.63
12	no	Kr	1993	51.56	41.68	63.76
13	yes	Ku	1993	31.26	24.47	39.93
14	no	Ku	1993	49.33	39.77	61.19
15	yes	Rd	1993	26.01	20.05	33.74
16	no	Rd	1993	41.04	31.76	53.03
17	yes	Ke	1994	20.80	16.08	26.90
18	no	Ke	1994	32.82	25.42	42.37
19	yes	Kr	1994	21.84	16.52	28.88
20	no	Kr	1994	34.47	26.69	44.52
21	yes	Ku	1994	20.90	15.78	27.69
22	no	Ku	1994	32.98	25.47	42.70
23	yes	Rd	1994	17.39	13.29	22.76
24	no	Rd	1994	27.44	21.00	35.84
25	yes	Ke	1995	27.17	21.05	35.07
26	no	Ke	1995	42.87	33.35	55.11
27	yes	Kr	1995	28.54	22.24	36.62
28	no	Kr	1995	45.03	36.11	56.15
29	yes	Ku	1995	27.30	21.22	35.13
30	no	Ku	1995	43.09	34.46	53.88
31	yes	Rd	1995	22.72	17.39	29.67
32	no	Rd	1995	35.84	27.55	46.63

This function does not require us to specify something like `tf = exp` because it automatically detects the link function and applies the appropriate function to produce the estimated expected response. The `glmint` function is particularly useful for making plots that include confidence intervals.


```
d <- cbind(d, trtools::glmint(m, newdata = d))
p <- ggplot(skylark, aes(x = count, y = spray)) +
  geom_pointrange(aes(x = fit, xmin = low, xmax = upp),
  shape = 21, fill = "white", data = d) +
  geom_point() + facet_grid(field + year ~ .) + theme_minimal() +
  labs(y = "Field Sprayed?", x = "Number of Skylark Fledglings")
```



Aflatoxicol and Liver Tumors in Trout

The data in the data frame `ex2116` in the **Sleuth3** package are from an experiment that investigated the relationship between aflatoxicol and liver tumors in trout. The figure below shows the proportion of trout in each tank that developed liver tumors as well as the dose of aflatoxicol to which the trout were exposed. Aflatoxicol is a metabolite of [Aflatoxin B1](#), a toxic by-product produced by a mold that infects some nuts and grains. Twenty tanks of rainbow trout embryos were exposed to one of five doses of aflatoxicol for one hour. The number of fish in each tank that developed liver tumors one year later was then observed. The plot below shows the data.

```
library(Sleuth3)
library(ggplot2)
p <- ggplot(ex2116, aes(x = Dose, y = Tumor/Total)) +
  geom_point(alpha = 0.5) + theme_minimal() + ylim(0, 1) +
  labs(x = "Dose (ppm)", y = "Proportion of Trout With Liver Tumors")
plot(p)
```



Note that `Tumor` is the number of trout in a tank that developed tumors, and `Total` is the number of trout in the tank. The goal here is to estimate the effect of aflatoxicol on the risk of liver tumors in trout. Here we will consider three different logistic regression models.

1. Estimating a logistic regression model for the probability of tumor development as a function of the dose of aflatoxicol as a quantitative explanatory variable. You should be able to replicate the following results.

```
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	-0.867	0.07673	-11.3	1.321e-29	-1.019	-0.7179
Dose	14.334	0.93695	15.3	7.838e-53	12.558	16.2346

Plot the model with the raw data, and estimate and interpret the odds ratio for the effect of increasing dose by 0.05 ppm.³

Solution: We can estimate the model as follows.


```
m <- glm(cbind(Tumor, Total - Tumor) ~ Dose, family = binomial, data = ex2116)
summary(m)$coefficients
```

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-0.867	0.07673	-11.3	1.321e-29
Dose	14.334	0.93695	15.3	7.838e-53

Here is a plot of the estimated model showing the probability of tumor development as a function of dose of aflatoxicol.

³Here e^{β_1} would be the odds ratio for the effect of increasing dose by 1 ppm. However that is probably not a realistic effect as it would be a relatively large increase in dose. The study only considered up to 0.25 ppm. Using `contrast` is convenient here to estimate the odds ratio for the effect of an arbitrary change in dose.

```
d <- data.frame(Dose = seq(0, 0.25, length = 100))
d$yhat <- predict(m, newdata = d, type = "response")
p <- ggplot(ex2116, aes(x = Dose, y = Tumor/Total)) +
  geom_point(alpha = 0.5) + theme_minimal() + ylim(0, 1) +
  geom_line(aes(y = yhat), data = d) +
  labs(x = "Dose (ppm)", y = "Proportion of Trout With Liver Tumors")
plot(p)
```



The plot suggests that the model does not fit the data well. But the odds ratio can be estimated as follows.

```
trtools::contrast(m,
  a = list(Dose = 0.1),
  b = list(Dose = 0.05), tf = exp)
```

```
estimate lower upper
2.048 1.868 2.245
```

```
pairs(emmeans(m, ~Dose, at = list(Dose = c(0.1, 0.05))),
  type = "response"), infer = TRUE)
```

contrast	odds.ratio	SE	df	asympt.LCL	asympt.UCL	null	z.ratio	p.value
Dose0.1 / Dose0.05	2.05	0.0959	Inf	1.87	2.24	1	15.298	<0.0001

Confidence level used: 0.95

Intervals are back-transformed from the log odds ratio scale

Tests are performed on the log odds ratio scale

The estimate odds ratio shows that the odds of tumor development increases by a factor of about 2.05 (i.e., about a 105% increase in the odds of tumor development) per 0.05 ppm increase in the dose of

aflatoxicol. Note that for this model the odds ratio is the same for *any* 0.05 ppm increase in the dose. For example, the same odds ratio would be found if dose was increased from 0.1 ppm to 0.15 ppm.

2. Estimate a logistic regression model like the one above but using the logarithm of the dose as the explanatory variable (i.e., apply a log transformation to dose). You should be able to replicate the following results.

```
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	4.163	0.20846	19.97	9.564e-89	3.763	4.581
log(Dose)	1.298	0.06434	20.17	1.628e-90	1.174	1.427

Plot the model with the raw data, and estimate and interpret the odds ratio for the effect of *doubling* the dose of aflatoxicol.


Solution: We can estimate the model as follows.

```
m <- glm(cbind(Tumor, Total-Tumor) ~ log(Dose), family = binomial, data = ex2116)
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	4.163	0.20846	19.97	9.564e-89	3.763	4.581
log(Dose)	1.298	0.06434	20.17	1.628e-90	1.174	1.427

Here is a plot of the estimated model showing the probability of tumor development as a function of dose of aflatoxicol.

```
d <- data.frame(Dose = seq(0, 0.25, length = 100))
d$yhat <- predict(m, newdata = d, type = "response")
p <- ggplot(ex2116, aes(x = Dose, y = Tumor/Total)) +
  geom_point(alpha = 0.5) + theme_minimal() + ylim(0, 1) +
  geom_line(aes(y = yhat), data = d) +
  labs(x = "Dose (ppm)", y = "Proportion of Trout With Liver Tumors")
plot(p)
```




This looks like an improvement, but a residual plot shows a trend which suggests that the model may still not have quite captured the relationship.

```

ex2116$yhat <- predict(m)
ex2116$residual <- rstudent(m)
p <- ggplot(ex2116, aes(x = yhat, y = residual)) + theme_minimal() +
  geom_point(alpha = 0.25) +
  labs(x = "Predicted Value (log odds scale)",
       y = "Studentized Residual")
plot(p)

```



The estimated odds ratio for the effect of doubling dose can be obtained as follows.

```
trtools::contrast(m,
  a = list(Dose = 0.2),
  b = list(Dose = 0.1), tf = exp)
```

```
estimate lower upper
2.459 2.253 2.684
```

```
pairs(emmeans(m, ~Dose, at = list(Dose = c(0.2, 0.1)),
  type = "response"), infer = TRUE)
```

contrast	odds.ratio	SE	df	asympt.LCL	asympt.UCL	null	z.ratio	p.value
Dose0.2 / Dose0.1	2.46	0.11	Inf	2.25	2.68	1	20.175	<0.0001

Confidence level used: 0.95

Intervals are back-transformed from the log odds ratio scale

Tests are performed on the log odds ratio scale

This odds ratio shows that doubling the dose of aflatoxin would increase the odds of tumor development by a factor of about 2.46 (i.e., about a 146% increase in the odds of tumor development).

3. Rather than trying to decide between using dose or some transformation of dose in the model, we can instead define dose as a 5-level factor. With this we do not need to assume a particular mathematical relationship between dose and the probability (or odds) of tumor development. But there are a couple of disadvantages. One is that inferences are limited to those dose values used in the study. Another is that it requires more parameters which can result in larger standard errors. There are two ways we could specify dose as a factor. One would be to create a new variable.

```
ex2116$Dosef <- factor(ex2116$Dose)
```

The levels of `Dosef` will be the original values of `Dose` but converted to strings, which we can see if we use the `levels` function.

```
levels(ex2116$Dosef)
```

```
[1] "0.01" "0.025" "0.05" "0.1" "0.25"
```

Another approach is to replace `Dose` in the model formula with `factor(Dose)`. Using this latter approach estimate a logistic regression model with dose as a categorical explanatory variable. Also estimate and interpret the odds ratios for the effect of a dose of 0.025 ppm versus 0.01 ppm, 0.05 ppm versus 0.01 ppm, 0.1 ppm versus 0.01 ppm, and 0.25 ppm versus 0.01 ppm.⁴

Solution: Here is how to estimate this model.

```
m <- glm(cbind(Tumor, Total-Tumor) ~ factor(Dose),
         family = binomial, data = ex2116)
cbind(summary(m)$coefficients, confint(m))
```

	Estimate	Std. Error	z value	Pr(> z)	2.5 %	97.5 %
(Intercept)	-2.556	0.2076	-12.310	8.049e-35	-2.988	-2.171
factor(Dose)0.025	2.073	0.2353	8.809	1.264e-18	1.628	2.553
factor(Dose)0.05	3.132	0.2354	13.306	2.130e-40	2.688	3.614
factor(Dose)0.1	3.890	0.2453	15.857	1.252e-56	3.427	4.391
factor(Dose)0.25	4.260	0.2566	16.605	6.436e-62	3.775	4.784

The odds ratios can be estimated as follows.

```
trtools::contrast(m, tf = exp,
                  a = list(Dose = c(0.025,0.05,0.1,0.25)),
                  b = list(Dose = 0.01))
```

estimate	lower	upper
7.945	5.01	12.60
22.920	14.45	36.36
48.909	30.24	79.10
70.840	42.84	117.13

```
contrast(emmeans(m, ~Dose, type = "response"), method = "trt.vs.ctrl",
         ref = 1, adjust = "none", infer = TRUE)
```

contrast	odds.ratio	SE	df	asympt.LCL	asympt.UCL	null	z.ratio	p.value
Dose0.025 / Dose0.01	7.94	1.87	Inf	5.01	12.6	1	8.809	<0.0001
Dose0.05 / Dose0.01	22.92	5.40	Inf	14.45	36.4	1	13.306	<0.0001
Dose0.1 / Dose0.01	48.91	12.00	Inf	30.24	79.1	1	15.857	<0.0001
Dose0.25 / Dose0.01	70.84	18.20	Inf	42.84	117.1	1	16.605	<0.0001

Confidence level used: 0.95

Intervals are back-transformed from the log odds ratio scale

Tests are performed on the log odds ratio scale

Note that in the `emmeans` package the `contrast` function is a bit different than the function of the same name in the `trtools` package, but there are some similarities in terms of what these functions are capable of doing. Here `method = "trt.vs.ctrl"` allows us to compare all but one of the levels with a “reference” level, which is specified by `ref = 1` meaning the first level as they are ordered (here, a dose of 0.01 ppm). The odds ratios show that the odds of tumor development at a dose of 0.025 ppm is about 7.94 times the odds at a dose of 0.01 ppm (i.e., about 694% higher), the odds of tumor

⁴Note that how you specify the levels of dose will depend on whether you created a new variable like `Dosef` or converted it to a factor within the model formula with `factor(Dose)`. For the latter you will need to specify dose as a *number* but if you created it to a new variable you will need to specify it as a *string* by enclosing it in quotes.

development at a dose of 0.05 ppm is about 22.92 times the odds at a dose of 0.01 ppm (i.e., about 2192% higher), the odds of tumor development at a dose of 0.1 ppm is about 48.91 times the odds at a dose of 0.01 ppm (i.e., about 4791% higher), and the odds of tumor development at a dose of 0.25 ppm is about 70.84 times the odds at a dose of 0.01 ppm (i.e., about 6984% higher).

4. Estimate the odds and probability of tumor development at each value of dose used in the study for any of the three models.

Solution: I will use the model from the previous problem for this. Using `contrast` the odds and probabilities can be estimated as follows.

```
trtools::contrast(m, a = list(Dose = c(0.01,0.025,0.05,0.1,0.25)),
  cnames = c(0.01,0.025,0.05,0.1,0.25), tf = exp) # odds
```

	estimate	lower	upper
0.01	0.07764	0.05168	0.1166
0.025	0.61682	0.49654	0.7662
0.05	1.77953	1.43188	2.2116
0.1	3.79730	2.93938	4.9056
0.25	5.50000	4.09298	7.3907

```
trtools::contrast(m, a = list(Dose = c(0.01,0.025,0.05,0.1,0.25)),
  cnames = c(0.01,0.025,0.05,0.1,0.25), tf = plogis) # probabilities
```

	estimate	lower	upper
0.01	0.07205	0.04914	0.1044
0.025	0.38150	0.33179	0.4338
0.05	0.64023	0.58880	0.6886
0.1	0.79155	0.74615	0.8307
0.25	0.84615	0.80365	0.8808

To estimate the odds using `emmeans` we need to use a “hack” that is not very intuitive.

```
emmeans(m, ~Dose, type = "response", tran = "log")
```

Dose	prob	SE	df	asympt.LCL	asympt.UCL
0.010	0.078	0.0161	Inf	0.052	0.117
0.025	0.617	0.0683	Inf	0.497	0.766
0.050	1.780	0.1970	Inf	1.432	2.212
0.100	3.797	0.4960	Inf	2.939	4.906
0.250	5.500	0.8290	Inf	4.093	7.391

Confidence level used: 0.95

Intervals are back-transformed from the log scale

Notice that somewhat confusingly the output still labels the estimates `prob` but these are odds as can be seen when comparing them with what was obtained using `contrast`. Estimated probabilities are simpler to obtain.

```
emmeans(m, ~Dose, type = "response")
```

Dose	prob	SE	df	asympt.LCL	asympt.UCL
0.010	0.072	0.0139	Inf	0.0491	0.104
0.025	0.382	0.0261	Inf	0.3318	0.434
0.050	0.640	0.0255	Inf	0.5888	0.689
0.100	0.791	0.0216	Inf	0.7462	0.831
0.250	0.846	0.0196	Inf	0.8037	0.881

Confidence level used: 0.95

Intervals are back-transformed from the logit scale

Here using `type = "response"` means that we want inferences on the scale of the response, which is a proportion, and the expected proportion is also the probability which is what we want.